

For older children

- Take deep breaths with your child to help “blow out” the pain.
- Point out interesting things in the room to help create distractions.
- Tell or read stories.
- Support your child if he or she cries. Never scold a child for not “being brave.”

Once your child has received all of the shots, be especially supportive. Hold, cuddle, and, for infants, breastfeed or offer a bottle. A soothing voice, combined with praise and hugs will help reassure your child that everything is ok.

After the Shots

Sometimes children experience mild reactions from vaccines, such as pain at the injection site, a rash or a fever. These reactions are normal and will soon go away. The following tips will help you identify and minimize mild side effects.

- Review any information your doctor gives you about the shots, especially the Vaccine Information Statements or other sheets that outline which side effects might be expected.
- Use a cool, wet cloth to reduce redness, soreness, and swelling in the place where the shot was given.
- Reduce any fever with a cool sponge bath. If your doctor approves, give non-aspirin pain reliever.
- Give your child lots of liquid. It’s normal for some children to eat less during the 24 hours after getting vaccines.
- Pay extra attention to your child for a few days. If you see something that concerns you, call your doctor.

*Remember to schedule your next visit!
Staying current with your child’s immunizations
provides the best protection against disease.*

*Take a moment to read the Vaccine Information
Sheet your health care professional gives you
during your visit. This sheet has helpful
information and describes possible side effects
your child may experience.*

